



# Derivation of the entropic formula for the statistical mechanics of space plasmas

George Livadiotis

Southwest Research Institute, Space Science & Engineering, San Antonio, TX, USA

5 *Correspondence to:* George Livadiotis (glivadiotis@swri.edu)

**Abstract.** Kappa distributions describe velocities and energies of plasma populations in space plasmas. The statistical origin of these distributions is the non-extensive statistical mechanics. Indeed, the kappa distribution is derived by maximizing the q-entropy of Tsallis under the constraints of canonical ensemble. However, there remains the question what is the physical origin of this entropic formulation. This paper shows that the q-entropy can be derived by adapting the additivity of energy and entropy.

10

## 1 Introduction

Space plasmas are collisionless and correlated particle systems characterized by a non-Maxwellian behavior, typically described by the formulations of kappa distributions. The origin of this vastly different statistical behavior between classical systems and space plasmas is the manifestation of correlations between the plasma particles. These systems are characterized by long-range interactions that induce correlations resulting to a collective behavior among particles (e.g., see Jund et al. 1995; Salazar & Toral 1999; Villain 2008; Tsallis 2009; Grassi 2010; Tirnakli & Borges 2016). The induction of any type of correlations among particles (more accurately, among particle energies or particle phase-space) departs the system from thermal equilibrium to be re-stabilized to other stationary states out of thermal equilibrium described by kappa distributions.

15

Kappa distributions describe numerous space plasma populations. Several examples are the following: (i) *inner heliosphere*, including solar wind (e.g., Maksimovic et al. 1997; Pierrard et al. 1999; Mann et al. 2002; Marsch 2006; Zouganelis 2008; Štverák et al. 2009; Livadiotis and McComas 2013a; Yoon 2014; Pierrard and Pieters 2015; Pavlos et al. 2016), solar spectra (e.g., Dzifčáková and Dudík 2013; Dzifčáková et al. 2015), solar corona (e.g., Owocki and Scudder 1983; Vocks et al. 2008; Lee et al. 2013; Cranmer 2014), solar energetic particles (e.g., Xiao et al. 2008; Laming et al. 2013), corotating interaction regions (e.g., Chotoo et al. 2000), and solar flares related (e.g., Mann et al. 2009; Livadiotis and McComas 2013b; Bian et al. 2014; Jeffrey et al. 2016); (ii) *planetary magnetospheres*, including magnetosheath (e.g., Formisano et al. 1973; Ogasawara et al. 2013), magnetopause (e.g., Ogasawara et al. 2015), magnetotail (e.g., Grabbe 2000), ring current (e.g., Pisarenko et al. 2002), plasma sheet (e.g., Christon 1987; Wang et al. 2003; Kletzing et al. 2003), magnetospheric substorms (e.g., Hapgood et al. 2011), Aurora (e.g., Ogasawara et al. 2017), magnetospheres of giant planets, such as Jovian (e.g., Collier and Hamilton 1995; Mauk et al. 2004), Saturnian (e.g., Dialynas et al. 2009; Livi et al.

25



2014; Carbary et al. 2014), Uranian (e.g., Mauk et al. 1987), Neptunian (Krimigis et al. 1989), magnetospheres of planetary moons, such as Io (e.g., Moncuquet et al. 2002) and Enceladus (e.g., Jurac et al. 2002), cometary magnetospheres (e.g., Broiles et al. 2016a; 2016b); (iii) *outer heliosphere and the inner heliosheath* (e.g., Decker and Krimigis 2003; Decker et al. 2005; Heerikhuisen et al. 2008; 2015; Zank et al. 2010; Livadiotis et al. 2011; 2012; 2013; Livadiotis and McComas 2011a; 5 2012; 2013c; Livadiotis 2014; 2016; Fuselier et al. 2014; Zirnstein and McComas 2015; Zank, 2015); (iv) *beyond the heliosphere*, including HII regions (e.g., Nicholls et al. 2012), planetary nebula (e.g., Nicholls et al. 2013; Zhang et al. 2014), and supernova magnetospheres (e.g., Raymond et al. 2010); and in cosmological scales (e.g., Hou et al. 2017); (iv) *other space plasma-related analyses* (e.g., Milovanov and Zelenyi 2000; Saito et al. 2000; Du 2004; Yoon et al. 2006; 2012; Raadu and Shafiq 2007; Livadiotis 2009; 2015a; 2015c; Tribeche et al. 2009; Hellberg et al. 2009; Livadiotis and McComas 10 2010b; 2014; Baluku et al. 2010; Le Roux et al. 2010; Eslami et al. 2011; Kourakis et al. 2012; Randol and Christian 2014; 2016; Varotsos et al. 2014; Fisk and Gloeckler 2014; Viñas et al. 2014, 2015; Ourabah et al. 2015; Dos Santos et al., 2016; Nicolaou and Livadiotis, 2016). (See also the book: Livadiotis 2017a, and references therein.)

Empirical kappa distributions have been introduced in mid-60's by Binsack (1966), Olbert (1968), and Vasyliūnas (1968), while their connection with statistical mechanics was shown and studied in detail in about half century later (see 15 Livadiotis and McComas 2009, and references therein). In particular, the statistical origin of these distributions is now widely accepted to be determined within the framework of non-extensive statistical mechanics (Tsallis 2009). This is a consistent generalization of the classical statistical mechanics, which is based on a mono-parametric ( $q$ -index) entropic formula (Tsallis 1988). The theoretical  $q$ -exponential distribution, which results from the maximization of entropy in the canonical ensemble, has the same formulation with the empirical kappa distribution; the two distributions are identical under 20 the transformation of their characteristic indices ( $q = 1 + 1/\kappa$ ).

Having attained a consistent connection of the mathematical model of kappa distributions with the physical means of entropy maximization does not precisely answer the main question regarding the origin of these distributions. We have shifted the modeling from the distributions to the entropic formulation. Therefore, we may understand now that the statistical origin of kappa distributions is given by the Tsallis entropy maximization in the canonical ensemble, but still, the origin of 25 this specific entropic formulation remains unknown.

Certainly, there are various mechanisms responsible for generating kappa distributions in space and other plasmas; for example, the presence of pickup ions (Livadiotis and McComas 2010a; 2011a) or weak turbulence (Yoon et al. 2012; Yoon 2014). Moreover, kappa distributions belong to the framework of non-extensive statistical mechanics. Thus, once a kappa distribution is generated and stabilized in a plasma population, the whole “tool package” of non-extensive statistical 30 mechanics is applicable for describing the statistical physics of this population; for instance, the entropy is given by the Tsallis formulation, while the temperature can be determined by the mean kinetic energy.

Here, we do not argue about whichever mechanisms generate kappa distributions in space plasmas, but for the physical reasons that these distributions sustain themselves in space plasmas once generated. The typical answer is that this is an effect of the presence and preservation of correlations in the collisionless environment that governs space plasmas. The



collisionless environment is preserving the energy leading to the additivity of energy: The energy of a multi-particle state is the sum of the energies of all the involved one-particle states. On the other hand, the preservation of local correlations among particles creates a conceptual separation of particles in correlation clusters. Debye spheres are correlation clusters that may include up to trillions of particles, since space plasmas are weakly coupled (Bryant 1996; Rubab and Murtaza 2006; Gougam and Tribeche 2011; Livadiotis and McComas 2014). This structure can lead to the additivity of entropy: The entropy of a multi-particle state is the sum of the entropies of all the involved one-particle states.

The purpose of this paper is to show that there is a deeper connection of Tsallis  $q$ -entropy and space plasmas: Namely, we will show that two simple first-principles such as the additive energy and additive entropy, which apply to plasma particle populations, are sufficient for indicating the specific formula of  $q$ -entropy (Figure 1). Therefore, the main objective of this work is to demonstrate the theory which determines that the entropic form given by the  $q$ -entropy formula proposed originally by Tsallis (1988) follows from certain assumptions regarding the (microscopic) state of the system. The importance of this discussion for the (space) plasma physics community resides mostly on the fact that the kappa velocity/energy distribution functions, ubiquitously observed in space and astrophysical environments, can be derived from the maximization of the  $q$ -entropy, under the constraints of a canonical ensemble.

In Section 2 we describe the physical motive of this paper in detail. In Section 3 we show in detail a similar property for both the entropic formalisms of BG and Tsallis: The entropy is non-additive in general for some arbitrary probability distribution; but it can become additive specifically for the canonical probability distribution (the one that maximizes the corresponding entropy). In Section 4 we show how we can determine the entropic formula appropriate for describing the plasma particle populations, simply by setting two first-principles properties, obvious for collisionless plasmas: energy and entropy are additive, at least macroscopically. Finally, Section 5 briefly summarizes the conclusions.




Figure 1: The infogram indicates the following triplet of concepts: (i) additive energy, (ii) additive entropy, and (iii) BG or Tsallis entropic formulation. Given any two out of the three features, the third can be derived. It is already known that BG or Tsallis entropic formulations can lead to additive entropy if the energy is also additive (red arrow 2). In the same way, it can be shown that these entropic formulations can lead to additive energy if the entropy is additive (purple arrow 1). The objective here is to show that the entropic formula can be derived from the additivity of energy and entropy (blue arrow 3).



## 2 Physical Motive

Classical Boltzmann Gibbs (BG) statistical mechanics characterizes systems with no correlations among particle velocities or energies. Therefore, the joint two-particle probability distribution can be expressed as the product of the one-particle identical independent discrete distributions; i.e., labelling the two particles with A and B,  $p_{ij}^{A+B} = p_i^A \cdot p_j^B$ . Hereafter, we consider a particle system described by a discrete energy spectrum  $\{\varepsilon_k\}_{k=1}^W$ , which is associated with a discrete probability distribution  $\{p_k\}_{k=1}^W$ . The same semantics is used when the system is separated in two subsystems A and B, where the two-particle distribution describes a two-particle state, with one particle at each subsystem. The logarithm of the probability is an additive function,  $\ln p_{ij}^{A+B} = \ln p_i^A + \ln p_j^B$ , from which we obtain the additivity of entropy,  $S^{A+B} = S^A + S^B$ . For special cases, however, where the independent relation does not apply,  $p_{ij}^{A+B} \neq p_i^A \cdot p_j^B$ , the entropy is non-additive,  $S^{A+B} \neq S^A + S^B$ . The logical reciprocate to the statements above is provided by the uniqueness theorem of Shannon (1948) and Khinchin (1960) that showed that under the assumption of the additivity of entropy (and other basic properties of entropy), the sufficient and necessary entropic form is given by the BG formula.

Non-extensive statistical mechanics characterizes systems with correlations among particles,  $p_{ij}^{A+B} \neq p_i^A \cdot p_j^B$ . For special systems, however, where the independent relation still applies, the entropy is non-additive,  $S^{A+B} \neq S^A + S^B$ ; in particular, a square, nonlinear term is added to the summation,  $S^{A+B} = S^A + S^B + (1-q)S^A S^B$  for some value of the entropic parameter  $q$  (where we set the Boltzmann constant  $k_B$  to 1). Note that the logical reciprocates exists also for this case, as shown by dos Santos (1997) and Abe (2000); namely, under the assumption of the mentioned non-additive property (and other basic properties of entropy), the sufficient and necessary entropic form is given by the  $q$ -entropic formula of Tsallis (1988).

Another property that is related with the additivity but is even more subtle and difficult to ascertain is the *extensivity* of the entropy. A non-additive entropy may be assumed to be also non-extensive, but it is the inverse assumption that is always correct, i.e., non-extensivity implies non-additivity). Nevertheless, certain correlations, expressed by the relation  $p_{ij}^{A+B} = g^{-1}[g(p_i^A) + g(p_j^B)]$  for some function  $g$ , can make the Tsallis entropy additive, and thus, recover its extensivity (e.g., Tsallis et al. 2005, Ruseckas 2015). In his book, C. Tsallis (2009) goes to great lengths to show that it is possible to find systems for which the BG entropy is *not* extensive. On the other hand, he argues that there are certain systems for which the entropic form can be extensive, for certain values of the entropic index  $q$ . In fact, he mentions in the preface that the term “nonextensive entropy” is somewhat incorrect in this sense, but it stuck for historical reasons.

The two statistical formalisms, classical BG and Tsallis non-extensive, have the common property that their entropy becomes additive for some specific function  $g$  in the relation  $p_{ij}^{A+B} = g^{-1}[g(p_i^A) + g(p_j^B)]$ , that is,  $g(x) \propto \ln(x)$  and



$g(x) \propto (x^{q-1} - 1)/(q - 1)$ , respectively; (the latter is related to the  $q$ -deformed logarithm; see: Silva et al. 1998; Yamano 2002).

It is important that the above probability relation is a characteristic feature of the canonical probability distribution in both the formalisms. In other words, the probability distribution that maximizes the BG entropy under the constraints of the canonical ensemble obeys to correlations expressed by  $g(x) \propto \ln(x)$  or  $p_{ij}^{A+B} = p_i^A \cdot p_j^B$ , which means zero correlation (due to the factorization of the exponentials, Livadiotis and McComas 2011b) that makes the entropy additive. Also, the probability distribution that maximizes the Tsallis entropy under the same constraints obeys to specific correlations expressed by  $g(x) \propto (x^{q-1} - 1)/(q - 1)$  or  $(p_{ij}^{A+B})^{q-1} = (p_i^A)^{q-1} + (p_j^B)^{q-1} - 1$ , which makes again the entropy additive. In Section 3 we show in detail this similar property of the two statistical formalisms.

Then, we may ask: Is the above described property of BG and Tsallis entropies a general feature of any physically meaningful entropic function? Or, can we reverse the question, and ask which specific entropic function obeys to the above properties? It will be really intriguing if we can determine the entropic formula appropriate for describing the plasma particle populations, simply by setting the following two first-principles properties: (1) additive energy, (2) additive entropy, i.e., the probability distribution derived by maximizing the entropy under the constraints of the canonical ensemble, makes the entropy additive. This will be the main purpose of this paper and will be examined in Section 4.

### 3 Canonical ensemble distributions with additive energy lead to additive entropy

#### 3.1 The Gibbs' path

The Gibbs' path (1902) for the maximization of the entropy  $S(p_1, p_2, \dots, p_W)$  under the constraints of canonical ensemble, i.e., (i) normalization  $1 = \sum_{k=1}^W p_k$ , and (ii) fixed internal energy  $U = \sum_{k=1}^W p_k \varepsilon_k$ , involves maximizing the functional

$$G(p_1, p_2, \dots, p_W) = S(p_1, p_2, \dots, p_W) + \lambda_1 \sum_{k=1}^W p_k + \lambda_2 \sum_{k=1}^W p_k \varepsilon_k \quad (1)$$

Next, we examine the BG and Tsallis entropic formulations.

#### 3.2 BG entropy

First, we start from the classical case of BG entropy

$$S(p_1, p_2, \dots, p_W) = - \sum_{k=1}^W p_k \ln(p_k) \quad (2)$$

where we ignored the Boltzmann constant  $k_B$  for simplicity. Then, setting  $(\partial/\partial p_j)G(p_1, p_2, \dots, p_W) = 0$  to



$$G(p_1, p_2, \dots, p_W) = -\sum_{k=1}^W p_k \ln(p_k) + \lambda_1 \sum_{k=1}^W p_k + \lambda_2 \sum_{k=1}^W p_k \varepsilon_k, \quad (3)$$

we find

$$p_j(\varepsilon_j) = \exp(\lambda_1 - 1) \cdot \exp(\lambda_2 \varepsilon_j). \quad (4)$$

We may write Eq.(4) in a logarithmic form,  $\ln p_j = \lambda_2 \varepsilon_j + \lambda_1 - 1$ . Then, we separate the particle system in two parts A and

5 B, so that each part is a new subsystem for which Eq.(4) holds:

$$\ln p_i^A = \lambda_2 \varepsilon_i^A + \lambda_1 - 1 \quad \text{and} \quad \ln p_j^B = \lambda_2 \varepsilon_j^B + \lambda_1 - 1. \quad (5)$$

The whole system is characterized by the joint probability,  $p_{ij}^{A+B}$ , meaning the probability of a particle in the subsystem A to reside at the state  $i$  and a particle in the subsystem B to reside at the state  $j$ . This is related with the energy  $\varepsilon_{ij}^{A+B}$  of the two-particle state,

$$10 \quad \ln p_{ij}^{A+B} = \lambda_2 \varepsilon_{ij}^{A+B} + \lambda_1 - 1. \quad (6)$$

Trivially, the energy of the two-particle state energy  $\varepsilon_{ij}^{A+B}$  equals the summation of the energy of each particle (since no interparticle force is considered), i.e., system's energy is additive:

$$\varepsilon_{ij}^{A+B} = \varepsilon_i^A + \varepsilon_j^B. \quad (7)$$

Hence, by eliminating energies from Eqs.(5,6), we find

$$15 \quad \ln p_{ij}^{A+B} + (\lambda_1 - 1) = \lambda_2 \varepsilon_i^A + (\lambda_1 - 1) + \lambda_2 \varepsilon_j^B + (\lambda_1 - 1) = \ln p_i^A + \ln p_j^B, \quad \text{or} \quad (8)$$

$$p_{ij}^{A+B} = p_i^A \cdot p_j^B \cdot e^{-(\lambda_1 - 1)}. \quad (9)$$

At this point we recall that the Lagrange multipliers,  $\lambda_1$  and  $\lambda_2$ , are related with the partition function  $Z = e^{-(\lambda_1 - 1)}$  and the inverse temperature  $\beta = -\lambda_2$ , respectively, and they are not necessarily equal for the two subsystems A and B, or the whole system A+B. Nevertheless, the logarithm of the partition function or  $(\lambda_1 - 1)$  is an extensive parameter, i.e.,

20  $(\lambda_1 - 1)^{A+B} = (\lambda_1 - 1)^A + (\lambda_1 - 1)^B$ , while the temperature is not an extensive parameter and can be considered the same

$\lambda_2^{A+B} = \lambda_2^A = \lambda_2^B$ . Then, instead of Eqs.(8,9), we obtain

$$\ln p_{ij}^{A+B} = \lambda_2 \varepsilon_{ij}^{A+B} + (\lambda_1 - 1)^{A+B} = \lambda_2 \varepsilon_i^A + (\lambda_1 - 1)^A + \lambda_2 \varepsilon_j^B + (\lambda_1 - 1)^B = \ln p_i^A + \ln p_j^B, \quad (10)$$

which clearly shows that the canonical probabilities are independent,

$$\ln(p_{ij}^{A+B}) = \ln(p_i^A) + \ln(p_j^B) \Rightarrow p_{ij}^{A+B} = p_i^A \cdot p_j^B. \quad (11)$$



Equation (9) indicates that the result in Eq.(11) can be obtained simply by setting  $\lambda_1=1$ . Certainly, this restricts the generality, but it can be used as a trick to simplify the calculations. Furthermore, we can easily obtain the additivity of entropy. Indeed, applying the operator  $\sum_{i=1}^W \sum_{j=1}^W p_{ij}^{A+B} \times$  on both sides of Eq.(11), we obtain

$$\begin{aligned} p_{ij}^{A+B} \ln p_{ij}^{A+B} &= p_{ij}^{A+B} \ln p_i^A + p_{ij}^{A+B} \ln p_j^B \\ \Rightarrow -\sum_{i=1}^W \sum_{j=1}^W p_{ij}^{A+B} \ln p_{ij}^{A+B} &= -\sum_{i=1}^W p_i^A \ln(p_i^A) - \sum_{j=1}^W p_j^B \ln(p_j^B) \end{aligned} \quad (12)$$

5 because  $\sum_{j=1}^W p_{ij}^{A+B} = p_i^A$ ,  $\sum_{i=1}^W p_{ij}^{A+B} = p_j^B$ . Hence, we arrive at the additivity of the entropy of the system to the entropies of the subsystems,

$$S^{A+B} = S^A + S^B \quad (13)$$

### 3.3 Tsallis entropy

Next, we continue with the Tsallis  $q$ -entropy,

$$10 \quad S(p_1, p_2, \dots, p_W) = \frac{1 - \phi(p_1, p_2, \dots, p_W)}{q-1} = \frac{1}{q-1} \cdot \sum_{k=1}^W (p_k - p_k^q) \quad (14)$$

where the argument is defined by

$$\phi(p_1, p_2, \dots, p_W) = \sum_{k=1}^W p_k^q \quad (15)$$

Again, the maximization of the entropy under the constraints of canonical ensemble involves maximizing the functional

$$15 \quad G(p_1, p_2, \dots, p_W) = \frac{1}{q-1} \cdot \sum_{k=1}^W (p_k - p_k^q) + \lambda_1 \sum_{k=1}^W p_k + \lambda_2 \sum_{k=1}^W p_k \varepsilon_k \quad (16)$$

Note that for simplicity we do not use the formulation of escort distributions (Beck and Schlogl 1993). The dyadic formalism of ordinary/escort distributions is of fundamental importance in the modern nonextensive statistical mechanics (Livadiotis 2017a; Chapter 1). It was shown that this dyadic formalism of distributions can be avoided in order to simplify the theory, but it leads to a dyadic formulation of entropy (Livadiotis 2017b).

Hence,  $(\partial/\partial p_j)G(p_1, p_2, \dots, p_W) = 0$ , gives

$$20 \quad p_j(\varepsilon_j) = \left[ 1 + (1-q^{-1}) \cdot (\lambda_1 - 1) \right]^{\frac{q^{-1}}{1-q^{-1}}} \cdot \left[ 1 + (1-q^{-1}) \cdot \frac{\lambda_2 \varepsilon_j}{1 + (1-q^{-1}) \cdot (\lambda_1 - 1)} \right]^{\frac{q^{-1}}{1-q^{-1}}}, \quad (17a)$$

or



$$p_j(\varepsilon_j) = \exp_{q^{-1}}^{q^{-1}}(\lambda_1 - 1) \cdot \exp_{q^{-1}}^{q^{-1}} \left[ \frac{\lambda_2 \varepsilon_j}{1_{q^{-1}}(\lambda_1 - 1)} \right], \quad (17b)$$

where reflects a generalization of Eq.(4). We used the  $Q$ -deformed exponential function, and its inverse, the  $Q$ -logarithm function (Silva et al. 1998; Yamano 2002), defined by

$$\exp_Q(x) = [1 + (1 - Q) \cdot x]_{+}^{-\frac{1}{Q-1}}, \quad \ln_Q(x) = \frac{1 - x^{1-Q}}{Q-1}. \quad (18a)$$

5 We also used the  $Q$ -deformed “unity function” (Livadiotis and McComas 2009), defined by

$$1_Q(x) = [1 + (1 - Q) \cdot x]_{+}, \quad (18b)$$

The subscript “+” in  $[...]_{+}$  denotes the cut-off condition, where  $\exp_Q(x)$  becomes zero if its base  $[...]$  is non-positive. Therefore, Eq.(17b) leads to

$$p_j^{q^{-1}} = 1 + (1 - q^{-1}) \cdot (\lambda_1 - 1) + (1 - q^{-1}) \cdot \lambda_2 \varepsilon_j, \quad (19)$$

$$10 \quad \frac{1 - p_j^{q^{-1}}}{q^{-1} - 1} = \ln_{q^{-1}}(p_j^q) = -q \cdot \ln_q(p_j^{-1}) = \lambda_2 \varepsilon_j + (\lambda_1 - 1), \quad (20)$$

Dividing again the whole system in two subsystems A and B, using the additivity of energy, and setting  $\lambda_1=1$ , the independence relation (11) is generalized to

$$\ln_q[(p_{ij}^{A+B})^{-1}] = \ln_q[(p_i^A)^{-1}] + \ln_q[(p_j^B)^{-1}] \Rightarrow (p_{ij}^{A+B})^{q^{-1}} = (p_i^A)^{q^{-1}} + (p_j^B)^{q^{-1}} - 1, \quad (21)$$

which is sometimes called  $q$ -independence relation (Umarov et al. 2008). Then, we apply the operator  $\sum_{i=1}^W \sum_{j=1}^W p_{ij}^{A+B} \times$ ,

$$15 \quad \sum_{i=1}^W \sum_{j=1}^W (p_{ij}^{A+B})^q = \sum_{i=1}^W (p_i^A)^q + \sum_{j=1}^W (p_j^B)^q - 1 \Rightarrow \phi^{A+B} = \phi^A + \phi^B - 1, \quad (22)$$

and using the entropic formula (14), we end up with the additivity of entropy, as shown in Eq.(13).

Note that the additivity leads to the extensivity: The additivity for some function  $f$  is expressed by  $f(A + B) = f(A) + f(B)$ , or considering  $N$  different subsystems,

$$f\left(\bigcup_{n=1}^N A_n\right) = \sum_{n=1}^N f(A_n), \quad (23a)$$

20 while the extensivity is expressed by

$$f\left(\bigcup_{n=1}^N A_0\right) = N \cdot f(A_0). \quad (23b)$$

Therefore, the canonical probability distribution, the one that maximizes the entropy under the constraints of canonical ensemble, makes the entropy additive (and therefore extensive) if the energy is additive. Several special conditions can simplify this result, e.g., constant Lagrange constraints with  $\lambda_1=1$ . This is true for both the entropic formulation of  
 25 classical BG and Tsallis nonextensive statistical mechanics.



Next, we will try to reverse the problem and seek to find the specific entropic formula, for which both the energy and entropy are additive.

#### 4 Additive energy and entropy leads to Tsallis entropic formalism

The general entropic form is still function of the probabilities,  $S = S(\{p_k\}_{k=1}^W)$ . Then, its derivative with respect to any of the probability components, let's say the  $i^{\text{th}}$ , is also a function of all of these component, i.e.,  $\partial S / \partial p_i = F_i(\{p_k\}_{k=1}^W)$ , for any  $i: 1, \dots, W$ . However, the 2<sup>nd</sup> constraint of the canonical ensemble connects the  $i^{\text{th}}$  entropic derivative to some function  $h_i$  of the  $i^{\text{th}}$  energy,  $\varepsilon_i$ , namely,  $\partial S / \partial p_i = h_i(\varepsilon_i)$ . On the other hand, the canonical probability distribution derived from the entropy maximization constitutes an expression of the  $i^{\text{th}}$  probability component with some function  $g$  of the  $i^{\text{th}}$  energy,  $p_i = g(\varepsilon_i)$ . Therefore, we conclude that  $\partial S / \partial p_i = F_i(p_i)$ , where  $F_i = h_i \circ g^{-1}$ ; in other words, the entropy can be factorized as a summation of functions of each probability component,  $S = \sum_{k=1}^W f_k(p_k)$ , where we set  $f_i(p_i) = \int F_i(p_i) dp_i$ . Finally, we consider that none of the probability components should have special effect on the entropy, i.e., the whole distribution “weights” in the same way, so that the entropic functional  $S = S(\{p_k\}_{k=1}^W)$  should be symmetric to any permutation of each components, e.g.,  $S = S(\dots, p_k, \dots, p_\ell, \dots) = S(\dots, p_\ell, \dots, p_k, \dots)$ . This, leads to  $f_k = f$ ; hence, considering (1) Entropy maximization, (2) No weighting, we obtain

$$S = \sum_{k=1}^W f(p_k) . \quad (24)$$

The maximization of entropy under the constraints of canonical ensemble, i.e.,  $1 = \sum_{k=1}^W p_k$  and  $U = \sum_{k=1}^W p_k \varepsilon_k$ , involves maximizing the functional  $G(\{p_k\}_{k=1}^W) = \sum_{k=1}^W f(p_k) + \lambda_1 \sum_{k=1}^W p_k + \lambda_2 \sum_{k=1}^W p_k \varepsilon_k$ . Hence, setting  $\partial G(\{p_k\}_{k=1}^W) / \partial p_i = 0$ , we obtain

$$F(p_i) + \lambda_1 + \lambda_2 \varepsilon_i = 0 , \text{ or } p_i(\varepsilon_i) = F^{-1}(-\lambda_1 - \lambda_2 \varepsilon_i) , \text{ with } F(x) \equiv f'(x) . \quad (25)$$

We now consider two systems A and B, with respective energy spectra  $\{\varepsilon_i^A\}_{i=1}^W$  and  $\{\varepsilon_j^B\}_{j=1}^W$ , associated with the discrete probability distributions  $\{p_i^A\}_{i=1}^W$  and  $\{p_j^B\}_{j=1}^W$ . The total system A+B has energy spectrum  $\{\varepsilon_{ij}^{A+B}\}_{i,j=1}^W$ , associated with the joint probability distribution  $\{p_{ij}^{A+B}\}_{i,j=1}^W$ . The probability distributions  $\{p_i^A\}_{i=1}^W$  and  $\{p_j^B\}_{j=1}^W$  are marginal of the joint distribution, i.e.,  $\sum_{j=1}^W p_{ij}^{A+B} = p_i^A$  and  $\sum_{i=1}^W p_{ij}^{A+B} = p_j^B$ . As we will find further below, the joint probability can be expressed as a function of the marginal probabilities,  $p_{ij}^{A+B} = h(p_i^A, p_j^B)$ . On the other hand, the relation between the joint



energies  $\varepsilon_{ij}^{A+B}$  is rather trivial to be derived: particles in A with energy  $\varepsilon_i^A$  and particles in B with energy  $\varepsilon_j^B$  ensemble the particles in A+B with energy  $\varepsilon_{ij}^{A+B} = \varepsilon_i^A + \varepsilon_j^B$ . Trivially, the same additivity holds for their mean values – the internal energies,

$$U^{A+B} = \sum_{i,j} p_{ij}^{A+B} \varepsilon_{ij}^{A+B} = \sum_{i,j} p_{ij}^{A+B} \varepsilon_i^A + \sum_{i,j} p_{ij}^{A+B} \varepsilon_j^B = \sum_i p_i^A \varepsilon_i^A + \sum_j p_j^B \varepsilon_j^B = U^A + U^B . \quad (26)$$

5 Now, the probability distributions are related to their energies, according to Eq.(7). According to Eq.(25), we have

$$F(p_i^A) + \lambda_1 + \lambda_2 \varepsilon_i^A = 0 , F(p_j^B) + \lambda_1 + \lambda_2 \varepsilon_j^B = 0 , F(p_{ij}^{A+B}) + \lambda_1 + \lambda_2 \varepsilon_{ij}^{A+B} = 0 , \quad (27)$$

and due to the additivity of energies, we obtain

$$F(p_{ij}^{A+B}) - \lambda_1 = F(p_i^A) + F(p_j^B) . \quad (28)$$

Again, the Lagrange constant,  $\lambda_1$  and  $\lambda_2$ , are considered to be constant. Setting  $\tilde{F} \equiv \frac{1}{-\lambda_1} F$ , Eq.(28) becomes

$$10 \quad [\tilde{F}(p_{ij}^{A+B}) - 1] = [\tilde{F}(p_i^A) - 1] + [\tilde{F}(p_j^B) - 1] , \text{ or,} \quad (29)$$

$$p_{ij}^{A+B} = h(p_i^A, p_j^B) , \text{ with } h(x, y) \equiv \tilde{F}^{-1}[\tilde{F}(x) + \tilde{F}(y) - 1] . \quad (30)$$

Then, we apply  $\sum_{i=1}^W \sum_{j=1}^W p_{ij}^{A+B}$  in both sides of Eq.(29),

$$\sum_{i,j=1}^W [\tilde{F}(p_{ij}^{A+B}) - 1] p_{ij}^{A+B} = \sum_{i=1}^W [\tilde{F}(p_i^A) - 1] \sum_{j=1}^W p_{ij}^{A+B} + \sum_{j=1}^W [\tilde{F}(p_j^B) - 1] \sum_{i=1}^W p_{ij}^{A+B} , \text{ or}$$

$$\sum_{i,j} [\tilde{F}(p_{ij}^{A+B}) - 1] p_{ij}^{A+B} = \sum_i [\tilde{F}(p_i^A) - 1] p_i^A + \sum_j [\tilde{F}(p_j^B) - 1] p_j^B . \quad (31)$$

15 (Note: The number of allowed states may be different for the two subsystems,  $W_A \neq W_B$ , but here it does not make any difference to consider  $W_A = W_B = W$ .)

We recall that  $\tilde{F}(x) \equiv \frac{1}{-\lambda_1} f'(x)$ , thus, we find

$$\sum_{i,j} \left[ \frac{1}{-\lambda_1} f'(p_{ij}^{A+B}) - 1 \right] p_{ij}^{A+B} = \sum_i \left[ \frac{1}{-\lambda_1} f'(p_i^A) - 1 \right] p_i^A + \sum_j \left[ \frac{1}{-\lambda_1} f'(p_j^B) - 1 \right] p_j^B . \quad (32)$$

We compare this property relation with the additivity of entropy

$$20 \quad S^{A+B} = \sum_{i,j} f(p_{ij}^{A+B}) = \sum_i f(p_i^A) + \sum_j f(p_j^B) = S^A + S^B . \quad (33)$$

The two functions  $f(x)$  and  $[\frac{1}{-\lambda_1} f'(x) - 1] \cdot x$  have the same additivity property. Therefore, one function  $f$  that can ensure for the additivity of entropy is the one that obeys to the proportionality,  $f(x) \propto [\frac{1}{-\lambda_1} f'(x) - 1] \cdot x$ , or to the differential equation



$$f(x) = c \cdot \left[ \frac{1}{-\lambda_1} f'(x) - 1 \right] \cdot x, \text{ or } f'(x) + \frac{\lambda_1}{c} \frac{1}{x} f(x) = -\lambda_1, \quad (34)$$

with solution

$$f(x) = \lambda_1 \cdot \frac{x - x^{\frac{\lambda_1}{c}}}{-\frac{\lambda_1}{c} - 1} + f(x=1) \cdot x^{\frac{\lambda_1}{c}}. \quad (35)$$

(Note: The selection of proportionality between the two functions  $f(x)$  and  $[\frac{1}{-\lambda_1} f'(x) - 1] \cdot x$  makes the derivation of Eq.(34)

5 a sufficient but not necessary condition. Other functionals may also exist; for example, a linear combination of the two mentioned functions.)

Then, we choose  $f(x=1) = 0$ , and we set  $q \equiv \frac{\lambda_1}{c}$ , where find

$$f(x) = \lambda_1 \cdot \frac{x - x^q}{q - 1}, \quad (36)$$

or, setting also  $\lambda_1 = 1$ ,

$$10 \quad f(x) = \frac{x - x^q}{q - 1}. \quad (37)$$

Therefore, the entropic function  $S = \sum_{k=1}^W f(p_k)$  becomes

$$S = \frac{1}{q - 1} \cdot \sum_{k=1}^W (p_k - p_k^q), \quad (38)$$

that is, the Tsallis entropic formulation that builds the nonextensive statistical mechanics.

## 5 Conclusions

15 The paper resolved a basic problem about the origin of the distributions and statistical mechanics applied in space plasmas. Kappa distributions, or combinations thereof, can describe the velocities and energies of the plasma populations in space plasmas. While these distributions were used since mid-60's for modeling space plasma datasets, their physical origin was remaining unknown. It was just the last decade that was completely understood that the statistical origin of these distributions is not the Boltzmann-Gibbs' classical statistical mechanics, but the Tsallis non-extensive statistical mechanics  
 20 (Livadiotis 2017a; Chapter 1). Indeed, the kappa distribution is the outcome of the maximization of the  $q$ -entropy of Tsallis under the constraints of canonical ensemble. Once this concept was understood by the science community, the next question was about the physical origin and reasoning of this entropic formula. This paper showed that the  $q$ -entropy, which is the entropic formula that maximized leads to the kappa distribution, can be derived under simple first-principles and conditions, namely, by considering that energy and entropy are both additive physical quantities.



**Acknowledgements:** The work was supported in part by the project NNX17AB74G of NASA's HGI Program.

## References

- Abe, S., 1997 A note on the q-deformation-theoretic aspect of the generalized entropies in nonextensive physics. *Physics Letters A* 224, 326 – 330.
- Abe, S., 2000 Axioms and uniqueness theorem for Tsallis entropy. *Physics Letters A* 271, 74–79.
- Baluku, T. K.; M. A. Hellberg, I. Kourakis, N. S. Saini, 2010 Dust ion acoustic solitons in a plasma with kappa-distributed electrons. *Phys. Plasmas* 17, 053702.
- Beck C. and Schlögl F. 1993 *Thermodynamics of Chaotic Systems*, Cambridge University Press.
- Bian, N., G.A. Emslie, D.J. Stackhouse, and E.P. Kontar 2014 The formation of a kappa-distribution accelerated electron populations in solar flares. *Astrophys. J.* 796, 142.
- Binsack, J. H. 1966 *Plasma studies with the IMP-2 satellite*, Ph.D. Thesis, MIT.
- Broiles, T.W., Livadiotis, G., Burch, J.L., Chae, K., Clark, G., Cravens, T.E., Davidson, R., Eriksson, A., Frahm, R.A., Fuselier, S.A., Goldstein, J., Goldstein, R., Henri, P., Madanian, H., Mandt, KE, Mokashi, P., Pollock, C., Rahmati, A., Samara, M., Schwartz, S.J. 2016a Characterizing cometary electrons with kappa distributions. *J. Geophys. Res.* 121, 7407-7422.
- Broiles, T.W., Burch, J.L., Chae, K., Clark, G., Cravens, T.E., Eriksson, A., Fuselier, S.A., Frahm, R.A., Gasc, S., Goldstein, R., Henri, P., Koenders, C., Livadiotis, G., Mandt, K.E., Mokashi, P., Nemeth, Z., Rubin, M. and Samara, M. 2016b Statistical analysis of suprathermal electron drivers at 67P/Churyumov-Gerasimenko. *MNRAS* 462, S312-S322.
- Carbary, J. F., M. Kane, B. H. Mauk and S. M. Krimigis 2014 Using the kappa function to investigate hot plasma in the magnetospheres of the giant planets. *J. Geophys. Res.* 119, 8426–8447.
- Chottoo, K., et al. 2000 The suprathermal seed population for corotaing interaction region ions at 1AU deduced from composition and spectra of H<sup>+</sup>, He<sup>++</sup>, and He<sup>+</sup> observed by Wind. *J. Geophys. Res.* 105, 23107–23122.
- Christon, S. P. 1987 A comparison of the Mercury and earth magnetospheres: electron measurements and substorm time scales. *Icarus* 71, 448-471.
- Collier, M.R., and D.C. Hamilton 1995 The relationship between kappa and temperature in the energetic ion spectra at Jupiter. *Geophys. Res. Lett.* 22, 303-306.
- Cranmer, S. R. 2014 Suprathermal electrons in the solar corona: Can nonlocal transport explain heliospheric charge states? *Astrophys. J. Lett.* 791, L31.
- Decker, R. B. and Krimigis, S. M. 2003 Voyager observations of low-energy ions during solar cycle 23. *Adv. Space Res.* 32, 597–602.
- Decker, R. B., et al. 2005 Voyager 1 in the foreshock, termination shock, and heliosheath. *Science* 309, 2020–2024.



- Dialynas, K., S. M. Krimigis, D. G. Mitchell, D. C. Hamilton, N. Krupp, and P. C. Brandt 2009 Energetic ion spectral characteristics in the Saturnian magnetosphere using Cassini/MIMI measurements. *J. Geophys. Res.* 114, A01212.
- Dos Santos, M. S., L. F. Ziebell, and R. Gaelzer 2016 Ion firehose instability in a dusty plasma considering product-bi-kappa distributions for the plasma particles. *Phys. Plasmas* 23, 013705.
- 5 Dos Santos, R. J. V., 1997 Generalization of Shannon's theorem for Tsallis entropy. *J. Math. Phys.* 38, 4104–4107.
- Du, J. 2004 The nonextensive parameter and Tsallis distribution for self-gravitating systems. *EPL* 67, 893–899.
- Dzifčáková, E., and J. Dudík 2013 H to Zn ionization equilibrium for the non-Maxwellian electron  $\kappa$ -distributions: Updated calculations, *Astrophys. J. Suppl. Ser.* 206, 6.
- Dzifčáková, E., J. Dudík, P. Kotrč, F. Fárník and A. Zemanová 2015 KAPPA: A package for synthesis of optically thin  
10 spectra for the non-Maxwellian  $\kappa$ -distributions based on the Chianti database. *Astrophys. J. Suppl. Ser.* 217, 14.
- Eslami, P., Mottaghizadeh, M., and Pakzad, H. R. 2011 Nonplanar dust acoustic solitary waves in dusty plasmas with ions and electrons following a q-nonextensive distribution. *Phys. Plasmas* 18, 102303.
- Fisk, L. A., and G. Gloeckler 2014 The case for a common spectrum of particles accelerated in the heliosphere: Observations and theory. *J. Geophys. Res.* 119, 8733.
- 15 Formisano, V., G. Moreno, F. Palmiotto, and P. C. Hedgecock 1973 Solar Wind Interaction with the Earth's Magnetic Field 1. Magnetosheath, *J. Geophys. Res.* 78, 3714–3730.
- Fuselier, S. A., Allegrini, F., Bzowski, M., Dayeh, M. A., Desai, M., Funsten, H. O., Galli, A., Heirtzler, D., Janzen, P., Kubiak, M. A., Kucharek, H., Lewis, W., Livadiotis, G., McComas, D. J., Möbius, E., Petrinec, S. M., Quinn, M., Schwadron, N., Sokół, J. M., Trattner, K. J., Wood, B. E., Wurz, P. 2014 Low energy neutral atoms from the heliosheath.  
20 *Astrophys. J.*, 784, 89.
- Gibbs, J.W., 1902, *Elementary Principles in Statistical Mechanics*, Scribner's sons, New York.
- Gougam, L. A. and Tribeche, M. 2011 Debye shielding in a nonextensive plasma. *Phys. Plasmas* 18, 062102.
- Grabbe, C. 2000 Generation of broadband electrostatic waves in Earth's magnetotail. *Phys. Rev. Lett.* 84, 3614.
- Grassi, A. 2010 A relationship between atomic correlation energy of neutral atoms and generalized entropy. *Int. J. Quantum*  
25 *Chem.* 111, 2390–2397.
- Hapgood, M., C. Perry, J. Davies and M. Denton 2011 The role of suprathermal particle measurements in CrossScale studies of collisionless plasma processes, *Planet. Space Sci.* 59, 618–629.
- Heerikhuisen, J., Pogorelov, N. V., Florinski, V., Zank, G. P. and le Roux, J. A. 2008 The effects of a k-distribution in the heliosheath on the global heliosphere and ENA flux at 1 AU. *Astrophys. J.* 682, 679–689.
- 30 Heerikhuisen, J., Zirnstein, E., Pogorelov, N. 2015  $\kappa$ -distributed protons in the solar wind and their charge-exchange coupling to energetic hydrogen. *J. Geophys. Res.* 120, 1516–1525.
- Hellberg, M.A., Mace, R.L., Baluku, T.K., Kourakis, I., Saini, N.S. 2009 Comment on "Mathematical and physical aspects of Kappa velocity distribution" [*Phys. Plasmas* 14, 110702 (2007)]. *Phys. Plasmas* 16, 094701.



- Hou, S. Q.; He, J. J.; Parikh, A.; Kahl, D.; Bertulani, C. A.; Kajino, T.; Mathews, G. J.; Zhao, G. 2017 Non-extensive Statistics to the Cosmological Lithium Problem. *Astrophys. J.* 834, 165.
- Jeffrey, N. L. S., Fletcher, L., and Labrosse, N. 2016 First evidence of non-Gaussian solar flare EUV spectral line profiles and accelerated non-thermal ion motion. *Astron. Astrophys.* 590, A99.
- 5 Jund, P., Kim, S. G., & Tsallis, C. 1995, Crossover from extensive to nonextensive behavior driven by long-range interactions. *Phys. Rev. B* 52, 50.
- Jurac, S., M. A. McGrath, R. E. Johnson, J. D. Richardson, V. M. Vasyliunas, and A. Eviatar 2002 Saturn: Search for a missing water source. *Geophys. Res. Lett.* 29, 2172.
- Khinchin, A. I. 1957 *Mathematical Foundations of Information Theory*. Dover Publications.
- 10 Kletzing, C. A., J. D. Scudder, E. E. Dors, and C. Curto 2003 Auroral source region: Plasma properties of the high latitude plasma sheet. *J. Geophys. Res.* 108, 1360.
- Kourakis, I., Sultana, S. Hellberg, M. A. 2012 Dynamical characteristics of solitary waves, shocks and envelope modes in kappa-distributed non-thermal plasmas: an overview. *Plasma Phys. Control. Fusion* 54, 124001.
- Krimigis, S. M., et al. 1989 Hot plasma and energetic particles in Neptune's magnetosphere. *Science* 246, 1483.
- 15 Laming, J. M., J. D. Moses, Y.-K. Ko, C. K. Ng, C. E. Rakowski, and A. J. Tylka (2013), On the remote detection of suprathermal ions in the solar corona and their role as seeds for solar energetic particle production. *Astrophys. J.* 770, 73.
- Le Roux, J. A.; G. M. Webb, A. Shalchi, G. P. Zank 2010 A generalized nonlinear guiding center theory for the collisionless anomalous perpendicular diffusion of cosmic rays. *Astrophys. J.* 716, 671-692.
- Lee, E., D. R. Williams, and G. Lapenta 2013 Spectroscopic indication of suprathermal ions in the solar corona, arXiv:1305.2939v.
- 20 Livadiotis, G. 2009 Approach on Tsallis statistical interpretation of hydrogen-atom by adopting the generalized radial distribution function. *J. Math. Chem.* 45, 930-939.
- Livadiotis, G. 2014 Lagrangian temperature: Derivation and physical meaning for systems described by kappa distributions. *Entropy* 16, 4290-4308.
- 25 Livadiotis, G. 2015a Statistical background and properties of kappa distributions in space plasmas. *J. Geophys. Res.* 120, 1607-1619.
- Livadiotis, G. 2015b Kappa distribution in the presence of a potential energy. *J. Geophys. Res.* 120, 880.
- Livadiotis, G. 2015c Kappa and q indices: Dependence on the degrees of freedom. *Entropy* 17, 2062.
- Livadiotis, G. 2016 Curie law for systems described by kappa distributions. *EPL* 113, 10003.
- 30 Livadiotis, G. 2017a *Kappa distribution: Theory & Applications in plasmas* Netherlands, UK, US: Elsevier.
- Livadiotis, G. 2017b On the simplification of statistical mechanics for space plasmas. *Entropy*, 19, 285.
- Livadiotis, G. and McComas, D.J. 2009 Beyond kappa distributions: Exploiting Tsallis statistical mechanics in space plasmas. *J. Geophys. Res.* 114, A11105.
- Livadiotis, G. and McComas, D.J. 2010a Exploring transitions of space plasmas out of equilibrium. *Astrophys. J.* 714, 971.



- Livadiotis, G. and McComas, D.J. 2010b Measure of the departure of the q-metastable stationary states from equilibrium. *Phys. Scr.* 82, 035003.
- Livadiotis, G. and McComas, D.J. 2011a The influence of pick-up ions on space plasma distributions. *Astrophys. J.* 738, 64.
- Livadiotis, G. and McComas, D.J. 2011b Invariant kappa distribution in space plasmas out of equilibrium. *Astrophys. J.* 741, 5 88.
- Livadiotis, G. and McComas, D.J. 2012 Non-equilibrium thermodynamic processes: Space plasmas and the inner heliosheath. *Astrophys. J.* 749, 11.
- Livadiotis, G. and McComas, D.J. 2013a Understanding kappa distributions: A toolbox for space science and astrophysics. *Space Sci. Rev.* 75, 183–214.
- 10 Livadiotis, G. and McComas, D.J. 2013b Evidence of large scale phase space quantization in plasmas. *Entropy* 15, 1118–1132.
- Livadiotis, G. and McComas, D.J. 2013c Near-equilibrium heliosphere – Far-equilibrium heliosheath. *AIP Conf. Proc.* 1539, 344–350.
- Livadiotis, G. and McComas, D.J. 2014 Electrostatic shielding in plasmas and the physical meaning of the Debye length. *J.* 15 *Plasma Phys.* 80, 341–378.
- Livadiotis, G., McComas, D.J., Dayeh, M.A., Funsten, H.O. and Schwadron, N.A. 2011 First sky map of the inner heliosheath temperature using IBEX spectra. *Astrophys. J.* 734, 1.
- Livadiotis, G., McComas, D. J., Randol, B., Möbius, E., Dayeh, M. A., Frisch, P. C., Funsten, H. O., Schwadron, N. A. and Zank, G. P. 2012 Pick-up ion distributions and their influence on ENA spectral curvature. *Astrophys. J.* 751, 64.
- 20 Livadiotis, G., McComas, D. J., Schwadron, N. A., Funsten, H. O. and Fuselier, S. A. 2013 Pressure of the proton plasma in the inner heliosheath. *Astrophys. J.* 762, 134.
- Livi, R., Goldstein, J., Burch, J. L., Crary, F., Rymer, A. M., Mitchell, D. G., and Persoon, A. M. 2014 Multi-instrument analysis of plasma parameters in Saturn’s equatorial, inner magnetosphere using corrections for spacecraft potential and penetrating background radiation. *J. Geophys. Res.* 119, 3683.
- 25 Maksimovic, M., V. Pierrard and J. Lemaire 1997 A kinetic model of the solar wind with Kappa distributions in the corona. *Astron. Astrophys.* 324, 725–734.
- Mann, G., H. T. Classen, E. Keppler, and E. C. Roelof 2002 On electron acceleration at CIR related shock waves. *Astron. Astrophys.* 391, 749–756.
- Mann, G., A. Warmuth, and H. Aurass 2009 Generation of highly energetic electrons at reconnection outflow shocks during solar flares. *Astron. Astrophys.* 494, 669–675.
- 30 Marsch, E. 2006 Kinetic physics of the solar corona and solar wind, *Living Rev. Solar Phys.* 3, 1.
- Mauk, B. H., S. M. Krimigis, E. P. Keath, A. F. Cheng, T. P. Armstrong, L. J. Lanzerotti, G. Gloeckler, and D. C. Hamilton 1987 The hot plasma and radiation environment of the Uranian magnetosphere. *J. Geophys. Res.* 92, 15283.



- Mauk, B. H., D. G. Mitchell, R. W. McEntire, C. P. Paranicas, E. C. Roelof, D. J. Williams, S. M. Krimigis, and A. Lagg 2004 Energetic ion characteristics and neutral gas interactions in Jupiter's magnetosphere. *J. Geophys. Res.* 109, A09S12.
- Milovanov A. V., and L. M. Zelenyi 2000 Functional background of the Tsallis entropy: "coarse-grained" systems and "kappa" distribution functions, *Nonlin. Processes Geophys.* 7, 211–221.
- 5 Moncuquet, M., F. Bagenal, and N. Meyer-Vernet 2002 Latitudinal structure of the outer Io plasma torus. *J. Geophys. Res.* 108, 1260.
- Nicholls, D. C., Dopita, M. A., Sutherland, R. S. 2012 Resolving the Electron Temperature Discrepancies in H II Regions and Planetary Nebulae:  $\kappa$ -distributed Electrons. *Astrophys. J.* 752, 148.
- Nicholls, D. C., Dopita, M. A., Sutherland, R. S., Kewley, L. J., Palay, E. 2013 Measuring nebular temperatures: the effect of new collision strengths with equilibrium and  $\kappa$ -distributed electron energies. *Astrophys. J. Supp.* 207, 21.
- 10 Nicolaou, G. and Livadiotis, G. 2016 Misestimation of temperature when applying Maxwellian distributions to space plasmas described by kappa distributions. *Astrophys. Space Sci* 361, 359.
- Ogasawara, K., Angelopoulos, V., Dayeh, M.A., Fuselier, S.A., Livadiotis, G., McComas, D.J., McFadden, J.P. 2013 Characterizing the dayside magnetosheath using ENAs: IBEX and THEMIS observations. *J. Geophys. Res.* 118, 3126-3137.
- 15 Ogasawara, K., Dayeh, M.A., Funsten, H.O., Fuselier, S.A., Livadiotis, G., and McComas, D.J. 2015 Interplanetary magnetic field dependence of the suprathermal energetic neutral atoms originated in subsolar magnetopause. *J. Geophys. Res.* 120, 964-972.
- Ogasawara, K., Livadiotis, G., Grubbs, G.A., Jahn, J.-M., Michell, R., Samara, M., Sharber, J. R., & Winningham, J. D., 2017 Properties of suprathermal electrons associated with discrete auroral arcs. *Geophys Res Let.* 44, 3475–3484.
- 20 Olbert, S. 1968 Summary of experimental results from M.I.T. detector on IMP-1, in *Physics of the Magnetosphere*, R. L. Carovillano, J. F. McClay, and H. R. Radoski (Eds), New York: Springer, 641.
- Ourabah, K., Ait Gougam, L., and Tribeche, M. 2015 Nonthermal and suprathermal distributions as a consequence of Superstatistics. *Phys. Rev. E* 91, 012133.
- Owocki, S.P., and J.D. Scudder 1983 The effect of a non-Maxwellian electron distribution on oxygen and iron ionization balances in the solar corona. *Astrophys. J.* 270, 758-768.
- 25 Pavlos, G.P.; O.E. Malandraki, E.G. Pavlos, A.C. Iliopoulos, L.P. Karakatsanis 2016 Non-extensive statistical analysis of magnetic field during the March 2012 ICME event using a multi-spacecraft approach. *Physica A* 464, 149–181.
- Pierrard, V. and Pieters, M. 2015 Coronal heating and solar wind acceleration for electrons, protons, and minor ions, obtained from kinetic models based on kappa distributions. *J. Geophys. Res.* 119, 9441.
- 30 Pierrard, V., M. Maksimovic, and J. Lemaire 1999 Electron velocity distribution function from the solar wind to the corona. *J. Geophys. Res.* 104, 17021-17032.
- Pisarenko, N. F., E.Yu. Budnik, Yu.I. Ermolaev, I.P. Kirpichev, V.N. Lutsenko, E.I. Morozova, and E.E. Antonova 2002 The ion differential spectra in outer boundary of the ring current: November 17, 1995 case study. *J. Atm. Solar-Terr. Phys.* 64, 573 – 583.



- Raadu, M. A., and Shafiq, M. 2007 Test charge response for a dusty plasma with both grain size distribution and dynamical charging. *Phys. Plasmas* 14, 012105.
- Randol, B.M. and Christian, E.R. 2014 Simulations of plasma obeying Coulomb's law and the formation of suprathermal ion tails in the solar wind. *J. Geophys. Res.* 119, 7025–7037.
- 5 Randol, B. M. and Christian, E.R. 2016 Coupling of charged particles via Coulombic interactions: Numerical simulations and resultant kappa-like velocity space distribution functions. *J. Geophys. Res.* 121, 1907–1919.
- Raymond, J. C., P. F. Winkler, W. P. Blair, J.-J. Lee, and S. Park 2010 Non-Maxwellian Ha profiles in Tycho's supernova remnant, *Astrophys. J.* 712, 901.
- Rubab, N. and Murtaza, G. 2006 Debye length in non-Maxwellian plasmas. *Phys. Scr.* 74, 145.
- 10 Ruseckas, J. 2015 Probabilistic model of N correlated binary random variables and non-extensive statistical mechanics. *Phys. Lett. A* 379, 654–659.
- Saito, S., F. R. E. Forme, S. C. Buchert, S. Nozawa, R. Fujii 2000 Effects of a kappa distribution function of electrons on incoherent scatter spectra, *Ann. Geophys.* 18, 1216-1223.
- Salazar, R., & Toral, R. 1999 Scaling Laws for a System with Long-Range Interactions within Tsallis Statistics. *Phys. Rev. Lett.* 83, 4233-4236.
- 15 Shannon, C. E., 1948 A mathematical theory of communication. *Bell System Techn. J.* 27, 379–423, 623–656.
- Silva, R., A. R. Plastino, and J. A. S. Lima 1998 A Maxwellian path to the q-nonextensive velocity distribution function. *Phys. Lett. A* 249, 401-408.
- Štverák, S., Maksimovic M., Travnicek P. M., Marsch E., Fazakerley A. N., and Scime E. E. 2009 Radial evolution of nonthermal electron populations in the low-latitude solar wind: Helios, Cluster, and Ulysses Observations. *J. Geophys. Res.* 114, A05104.
- 20 Tirnakli, U., & Borges, E. P. 2016, The standard map: From Boltzmann-Gibbs statistics to Tsallis statistics. *Scientific Reports* 6, 23644.
- Tribeche, M., Mayout, S. and Amour, R. 2009 Effect of ion suprathermality on arbitrary amplitude dust acoustic waves in a charge varying dusty plasma. *Phys. Plasmas* 16, 043706.
- 25 Tsallis, C. 1988 Possible generalization of Boltzmann-Gibbs statistics. *J. Stat. Phys.* 52, 479–487.
- Tsallis, C. 2009 *Introduction to Nonextensive Statistical Mechanics*, New York: Springer.
- Tsallis, C., Gell-Mann, M. and Sato, Y. 2005 Asymptotically scale-invariant occupancy of phase space makes the entropy Sq extensive. *PNAS* 102, 15377–15382.
- 30 Umarov, S., Tsallis, C. and Steinberg, S. 2008 On a q-central limit theorem consistent with nonextensive statistical mechanics. *Milan J. Math.* 76, 307
- Varotsos, P.A., Sarlis, N.V., Skordas, E.S., 2014 Study of the temporal correlations in the magnitude time series before major earthquakes in Japan. *J. Geophys. Res.* 119, 9192–9206



- Vasyliūnas, V. M. 1968 A survey of low-energy electrons in the evening sector of the magnetosphere with OGO 1 and OGO 3. *J. Geophys. Res.* 73, 2839–2884.
- Villain, J. 2008, On the long-range interactions and non-extensive systems. *Scientifica Acta* 2, 93-99.
- Viñas, A. F., P. S. Moya, R. Navarro, and J. A. Araneda 2014 The role of higher-order modes on the electromagnetic whistler-cyclotron wave fluctuations of thermal and non-thermal plasmas, *Phys. Plasmas* 21, 012902.
- Viñas, A. F., P. S. Moya, R. E. Navarro, J. A. Valdivia, J. A. Araneda, and V. Muñoz 2015 Electromagnetic fluctuations of the whistler-cyclotron and firehose instabilities in a Maxwellian and Tsallis-kappa-like plasma, *J. Geophys. Res.* 120, 3307-3317.
- Vocks, C., G. Mann, and G. Rausche 2008 Formation of suprathermal electron distributions in the quiet solar corona. *Astron. Astrophys.* 480, 527–536.
- Wang, C.-P., L. R. Lyons, M. W. Chen, R. A. Wolf, and F. R. Toffoletto 2003 Modeling the inner plasma sheet protons and magnetic field under enhanced convection. *J. Geophys. Res.* 108, 1074.
- Xiao, F., C. Shen, Y. Wang, H. Zheng, and S. Whang 2008 Energetic electron distributions fitted with a kappa-type function at geosynchronous orbit. *J. Geophys. Res.* 113, A05203.
- Yamano, T. 2002 Some properties of q-logarithmic and q-exponential functions in Tsallis statistics. *Physica A* 305, 486–496.
- Yoon, P. H. 2014 Electron kappa distribution and quasi-thermal noise. *J. Geophys. Res.* 119, 7074.
- Yoon, P. H.; T. Rhee, C. M. Ryu, 2006 Self-consistent formation of electron  $\kappa$  distribution: 1. Theory, *J. Geophys. Res.* 111, A09106.
- Yoon, P. H.; L.F. Ziebell, R. Gaelzer, R.P. Lin, L. Wang, 2012 Langmuir turbulence and suprathermal electrons, *Space Sci. Rev.* 173, 459–489.
- Zank, G. P.; J. Heerikhuisen, N. V. Pogorelov, R. Burrows, D. J. McComas 2010 Microstructure of the heliospheric termination shock: implications for energetic neutral atom observations. *Astrophys. J.* 708, 1092.
- Zank, G. P. 2015 Faltering steps into the galaxy: The boundary regions of the heliosphere. *Ann. Rev. Astron. Astrophys.* 53, 449.
- Zhang, Y., Liu, X.-W. and Zhang B. 2014 H-I free-bound emission of planetary nebulae with large abundance discrepancies: Two-component models vs.  $\kappa$ -distributed electrons. *Astrophys. J.* 780, 93.
- Zirnstein, E. J. and McComas, D. J. 2015 Using kappa functions to characterize outer heliosphere proton distributions in the presence of charge-exchange. *Astrophys. J.* 815, 31.
- Zouganelis, I. 2008 Measuring suprathermal electron parameters in space plasmas: Implementation of the quasi-thermal noise spectroscopy with kappa distributions using in situ Ulysses/URAP radio measurements in the solar wind. *J. Geophys. Res.* 113, A08111.